arXiv:quant-ph/9909019v1 6 Sep 1999

Comparisons of spectra determined using detector atoms and spatial correlation functions

M. Havukainen
Helsinki Institute of Physics, PL 9, FIN-00014 Helsingin yliopisto, Finland

(February 1, 2008)

We show how two level atoms can be used to determine the local time dependent spectrum. The method is applied to a one dimensional cavity. The spectrum obtained is compared with the mode spectrum determined using spatially filtered second order correlation functions. The spectra obtained using two level atoms give identical results with the mode spectrum. One benefit of the method is that only one time averages are needed. It is also more closely related to a realistic measurement scheme than any other definition of a time dependent spectrum.

I. INTRODUCTION

It has turned out to be difficult to give a universally accepted definition of the time dependent spectrum. There are several definitions [1–3] which have not been popular in optics. A definition, which is connected to a realistic spectrum measurement was given by Eberly and Wódkiewicz [4]. Their, 'physical spectrum', has become a kind of canonical definition for a time dependent spectrum in quantum optics. All definitions mentioned above are based on integrals, typically Fourier transforms, over different two time averages of classical stochastic variables or quantum mechanical operators.

In our earlier paper [5], a totally new approach was introduced. Instead of calculating two time averages the radiation is guided to a group of two level atoms. All the atoms have equal, very small decay constants, but the resonance frequencies are all different. The spectrum measured by the atoms can be read from the excitation probabilities of the atoms. We have studied the spectrum of resonance fluorescence radiation from a laser driven three level atom [5]. The method was shown to give exactly the same result as the 'physical spectrum' defined by Eberly and Wódkiewicz. In order to calculate the two time averages needed to determine the 'physical spectrum', the Quantum Regression Theorem (QRT) [6] must be used. It has been shown that there are systems where QRT cannot be used i.e. the two time averages it gives are incorrect [7]. This is the case if the interaction between the matter, which emits the radiation, and the radiation field is strong. Interaction can be strong e.g. in microcavities. In the method of analyzer atoms, two time averages are not needed. The method can be used also in situations where the QRT is known to give incorrect results. It can be said that it is closer to a realistic spectrum measurement than any other definition.

In our earlier paper, the three level atom was coupled to the analyzer atoms using cascaded master equations [8,9]. Thus the quantum mechanical state of the field was not made available. In this paper we connect the method of analyzer atoms to our cavity QED simulations in one dimension. In these simulations there are two level atoms inside a one dimensional cavity. The field is quantized using canonical quantization. The state vector is restricted to have only one excitation i.e. the field strength is very low. The model was introduced by V. Bužek et. al. [10,11] and has been used in several different studies in one [12] and two dimensions [13]. The quantum mechanical state of the field is available so the spectrum of the radiation can be read directly from the excitations of the modes. The spatial dependence of the spectrum is obtained by calculating the second order correlation function $g(r_1, r_2)$ and using appropriate filter functions. The typical situation in the simulations is that the initial field intensity is split into left and right propagating parts using two level atoms. The spectra of both parts is determined using both analyzer atoms and filtered correlation functions. In all cases the two different methods give identical results.

In Sec. II we specify the model used in the simulations. Spatial correlation functions and their relation to the spectrum is explained in Sec. III. In Sec. IV we present the analyzer atom method for spectrum measurements. In Sec. IV we show several results of our simulations and finally in Sec. VI we present our conclusions.

II. HAMILTONIANS AND STATE VECTORS

A. Hamiltonians

In our simulations the mode functions in the one dimensional cavity ($0 \leq r \leq L$) are

$$G_n(r) = \sin(k_n r), \quad k_n = n\frac{\pi}{L}, \quad n = 1, 2, 3... \tag{1}$$

All mode functions are zero at the edges of the cavity, $r = 0$ and $r = L$. The electric and magnetic field operators can be expanded using the mode functions (1)

$$\hat{E} = \sum_{n=1}^{\infty} \left(\frac{\hbar\omega_n}{\epsilon_0 L}\right)^{1/2} \sin(k_n r)(\hat{a}_n + \hat{a}_n^\dagger) \tag{2}$$

$$\hat{B} = i\sum_{n=1}^{\infty} \left(\frac{\hbar\omega_n \mu_0}{L}\right)^{1/2} \sin(k_n r)(\hat{a}_n - \hat{a}_n^\dagger),$$

where $\hat{a}$ and $\hat{a}^\dagger$ are annihilation and creation operators. The operators satisfy the usual canonical commutation relations

$$[\hat{a}_n, \hat{a}_m^\dagger] = \delta_{nm}, \tag{3}$$
$$[\hat{a}_n, \hat{a}_m] = [\hat{a}_n^\dagger, \hat{a}_m^\dagger] = 0. \tag{4}$$

For the energy density operator we get using the expansions (2)

$$\begin{aligned}\hat{H}(r) &= \frac{1}{2}\epsilon_0 \hat{E}(r)^2 + \frac{1}{2\mu_0}\hat{B}(r)^2 \qquad (5)\\ &= \frac{2\hbar}{L}\sum_{n=1}^{\infty}\sum_{n'=1}^{\infty}\sqrt{\omega_n\omega_{n'}}\sin(k_n r)\sin(k_{n'} r)(\hat{a}_{n'}^\dagger\hat{a}_n + \frac{1}{2}).\end{aligned}$$

Integration of $\hat{H}(r)$ over the whole cavity and use of the orthogonal integral

$$\int_0^L \sin(k_n r)\sin(k_{n'} r)dr = \frac{L}{2}\delta_{nm}, \quad n, m \geq 1 \tag{6}$$

gives the familiar field Hamiltonian

$$\hat{H}_F = \sum_{n=1}^{\infty}\hbar\omega_n(\hat{a}_n^\dagger\hat{a}_n + \frac{1}{2}). \tag{7}$$

At fixed positions inside the cavity there are N_A two level atoms with resonance frequencies ω_j and dipole constants D_j. The atomic Hamiltonian is the sum over all one atom Hamiltonians

$$\hat{H}_A = \sum_{j=1}^{N_A}\hbar\omega_j\hat{\sigma}_z^j, \tag{8}$$

where $\hat{\sigma_z}$ is a Pauli spin matrix. The radiation field and the atoms are coupled through dipole coupling. The dipole operator of the j:th atom is

$$\hat{D}_j = (D_j\hat{\sigma}_+^j + D_j^*\hat{\sigma}_-^j). \tag{9}$$

For the interaction Hamiltonian we get

$$\hat{H}_I = -\sum_{j=1}^{N_A}\hat{D}_j\cdot\hat{E}(r_j), \tag{10}$$

where $\hat{E}(r_j)$ is the electric field operator (2) at the atomic position r_j. Using the expansion (2) for the field operator and (9) for the dipole operator we get

$$\hat{H}_I = -\sum_{j=1}^{N_A}\sum_{n=1}^{\infty}\left(\frac{\hbar\omega_n}{\epsilon_0 L}\right)^{1/2}\sin(k_n r)(D_j\hat{\sigma}_+^j\hat{a}_n + D_j^*\hat{\sigma}_-^j\hat{a}_n^\dagger). \tag{11}$$

The terms $\hat{\sigma}_+^j\hat{a}_j^\dagger$ and $\hat{\sigma}_-^j\hat{a}_j$, which do not affect the time evolution significantly, have been neglected. This approximation is called the Rotating Wave approximation (RWA). The Hamiltonian (11) has the familiar Jaynes-Cummings form. The total Hamiltonian is the sum of field, atomic and interaction Hamiltonians:

$$\hat{H} = \hat{H}_F + \hat{H}_A + \hat{H}_I. \tag{12}$$

B. State vectors

The general state vector is restricted to have only one excitation. The most general state vector has the form

$$\begin{aligned}|\Psi\rangle &= \sum_k\left(c_k|1\rangle_k\prod_{k'\neq k}|0\rangle_{k'}\right)\otimes\prod_{j=1}^{N_A}|0\rangle_j\\ &\quad+\prod_k|0\rangle_k\otimes\sum_{j=1}^{N_A}\left(c_j|1\rangle_j\prod_{j'=1,j'\neq j}^{N_A}|0\rangle_{j'}\right) \qquad (13)\\ &\equiv\sum_k c_k|1_k,0\rangle + \sum_{j=1}^{N_A}c_j|0,1_j\rangle.\end{aligned}$$

In the first sum, one of the field basis vectors have one excitation and in the second one, the excitation is in the atomic basis functions. If the initial state vector has only one excitation it does not obtain more excitations because the RWA was used in the interaction Hamiltonian. The state vector at all times is given by equation (13).

The state vector used as the initial state in our simulations is of the form

$$|\Psi\rangle = \sum_k(2\pi\sigma_k^2)^{(-1/4)}\exp\left(-ikr_0 - \frac{(k-k_0)^2}{4\sigma_k^2}\right)|1_k, 0\rangle. \tag{14}$$

The mode distribution $|c_k|^2$ is a Gaussian centered at $k = k_0$ with a variance σ_k^2. The phase factor e^{-ikr_0} is important. It guarantees that the energy density distribution is also a Gaussian centered at r_0.

III. CORRELATION FUNCTIONS AND THE MODE SPECTRUM

A. Correlation functions

The exact quantum mechanical state of the field is determined uniquely if all correlation functions of the operators $\hat{E}(r)$ and $\hat{B}(r)$ are known. In our simulations, because of the special form of the state vector (13) the second order correlation function determines the field state uniquely. In the following we denote $\epsilon_0 = \mu_0 = 1$. Using the expansions (2) we get for the normally ordered correlation function

$$\begin{aligned}
&\langle\Psi(t)|:\hat{E}(r_1)\hat{E}(r_2):|\Psi(t)\rangle \\
&\quad=\sum_{n=1}^{\infty}\sum_{m=1}^{\infty}\frac{1}{L}\sqrt{\omega_n\omega_m}\sin(k_nr_1)\sin(k_mr_2)\langle\Psi(t)|\hat{a}_n^{\dagger}\hat{a}_m+\hat{a}_m^{\dagger}\hat{a}_n|\Psi(t)\rangle \qquad (15)\\
&\quad=\sum_{n=1}^{\infty}\sum_{m=1}^{\infty}\frac{1}{L}\sqrt{\omega_n\omega_m}\sin(k_nr_1)\sin(k_mr_2)(c_n^*(t)c_m(t)+c_m^*(t)c_n(t))\\
&\quad=T^*(r_1,t)T(r_2,t)+T(r_1,t)T^*(r_2,t)
\end{aligned}$$

where

$$T(r,t)=\sum_{n=1}^{\infty}\sqrt{\frac{\omega_n}{L}}\sin(k_nr)c_n(t). \qquad (16)$$

A similar calculation gives

$$\begin{aligned}
&\langle\Psi(t)|:\hat{B}(r_1)\hat{B}(r_2):|\Psi(t)\rangle \qquad (17)\\
&=T^*(r_1,t)T(r_2,t)-T^*(r_2,t)T(r_1,t).
\end{aligned}$$

It is possible to invert the formula (16) and get the mode coefficients in Eq. (13). Multiplying both sides by $\sin(k_mr)$ and using the orthogonality integral (6) we get

$$c_m(t)=\frac{2}{\sqrt{\omega_mL}}\int_0^L\sin(k_mr)T(r,t)dr. \qquad (18)$$

So if $T(r,t)$ is known the mode coefficients can be calculated. It is also possible to calculate the coefficients c_n from the correlation functions (15) and (17). Let us define

$$\begin{aligned}
&W(r_1,r_2)\\
&=\langle\Psi(t)|\left(:\hat{E}(r_1)\hat{E}(r_2):+:\hat{B}(r_1)\hat{B}(r_2):\right)|\Psi(t)\rangle\\
&=T^*(r_1)T(r_2) \qquad (19)
\end{aligned}$$

Integrating both sides by $\int_o^L\sin(k_1r)\sin(k_pr)dr$ we get

$$c_1^*c_p=\frac{4}{L\sqrt{\omega_1\omega_p}}\int_0^L dr_1\int_0^L dr_2\sin(k_1r_1)\sin(k_pr_2)W(r_1,r_2). \qquad (20)$$

This gives equations for each c_p. For $p=1$ we get $|c_1|^2$. We can choose c_1 to be real and determine c_1. The rest of the coefficients are now determined uniquely. Normalization can be used as a check of the calculation.

B. A local mode spectrum

The mode spectrum $|c_k|^2$ gives the time dependent spectrum of the radiation in the whole cavity. In order to get a spectrum of radiation in some part of the cavity only we use filtered correlation functions. The correlation function (19) is replaced by a filtered correlation function

$$W_F(r_1,r_2;r_0)=g^F(r_1,r_2,r_0)W(r_1,r_2), \qquad (21)$$

where the filter function is a real-valued window function centered at r_0. The method is similar to the Windowed Fourier Transform (WFT), Ref. [14,15], which is used to determine the time dependent frequency distribution of a time dependent signal. The mode spectrum is calculated using a filtered correlation function (21) in equation (20). The spectrum obtained is the mode spectrum of the radiation in the part of the cavity where the filter function is nonzero.

A typical example of the filter function of one variable is a Gaussian filter

$$g^F(r;r_0)=(2\pi\sigma_r^2)^{-1/2}\exp\left(-\frac{(r-r_0)^2}{2\sigma_r^2}\right), \qquad (22)$$

where σ_r^2 is the variance of the window. Two dimensional filter is a product of two one dimensional ones. Another filter used in our simulations is a constant filter which is unity in some region and zero everywhere else

$$r=\begin{cases}1 & r_{min}\le r\le r_{max}\\ 0 & r\le r_{min},\quad r\ge r_{max}\end{cases}. \qquad (23)$$

The general behavior of the spectrum with all filter functions is such that the broader the filter function, the smaller details can be seen in the spectrum.

IV. TRADITIONAL DEFINITIONS OF THE SPECTRUM AND ANALYZER ATOMS

A. Traditional definitions of the spectrum

In the previous section the spectrum was calculated using '*spatial correlation functions at a specific time value*'. The traditional method to get the spectrum is to calculate the Fourier transform of two time averages of field operators, i.e. use '*time correlation functions at a specific point*'. For a stationary field the spectrum can be defined as

$$S(\omega)=\int_{-\infty}^{\infty}d\tau\langle\hat{E}(t)\hat{E}(t+\tau)\rangle e^{i\omega\tau}. \qquad (24)$$

This definition is valid only for stationary fields. There are many generalizations of a Fourier spectrum for non-stationary fields [1–3]. A method which takes also the measurement scheme into account was presented by Eberly and Wódkiewicz [4]. They define a 'physical spectrum' to be a double Fourier transform of two time averages multiplied by filter functions

$$S_{phys}(t,\omega_f,\Gamma_f)=\Gamma_f^2\int\limits_0^\infty d\tau_1\int\limits_0^\infty d\tau_2\exp(-(\Gamma_f-i\omega_f)\tau_1)\exp(-(\Gamma_f+i\omega_f)\tau_2)\langle\hat{E}^*(t-\tau_1)\hat{E}(t-\tau_2)\rangle. \quad (25)$$

The frequency filters are placed in front of the photodetector. They allow only the radiation with a certain frequency to pass to the detector. The spectrum is obtained from the relative intensities measured by the detector when the filters are tuned to different frequencies. The spectrum obtained is time dependent and the definition incorporates the measurement scheme. Our method to include time dependence to the spectrum is analogous to the method shown in the previous section. The definition (25) has '*time filtered*' correlation functions whereas in the last section '*spatially filtered*' correlation functions were used.

In many problems of quantum optics the radiation field is traced out from the Hilbert space. As a result the time evolution of the system is described by a master equation, which does not have information about the field degrees of freedom. Using this approach it is not possible to calculate the spatial correlation functions; then the definitions with time correlation functions like (24) and (25) are the only useful definitions.

B. Analyzer atoms

In our earlier paper [5], a totally different approach to the time dependent spectrum was introduced. The radiation from the system of interest is guided to a group of N two level atoms which all have the same very small decay constant Γ. The resonance frequencies of the atoms are all different

$$\omega_n=n\cdot\Delta\omega,\qquad \Delta\omega=\frac{\omega_{max}-\omega_{min}}{N-1},\qquad n=1,2...N. \quad (26)$$

Initially all atoms are in the ground state. The incoming radiation excites the atoms. Because the decay constants are equal and very small, the excitation probabilities of the atoms are directly proportional to the intensity of the incoming radiation at the resonance frequency. The normalized excitation as a function of resonance frequency of the atoms gives the time dependent spectrum of the radiation. A detailed description of the method and a comparison with a spectrum (25) calculated using two time averages can be found in Ref. [5]. In these simulations cascaded master equations were used to describe a three level atom and a group of analyzer atoms. The two spectra were shown to give identical results for the resonance fluorescence radiation of a three level atom.

The analyzer atom method can be used in our cavity simulations. Now the atoms with frequencies given by Eq. (26) and very small decay constants are located at some point inside the cavity. As in the case of master equations, the excitations should give the local time dependent spectrum of the radiation which passes them. In the next section the spectrum measured using detector atoms is compared with the mode spectrum calculated using filtered spatial correlation functions.

V. SIMULATIONS

In this section we show results of spectrum measurements in three different cases. In all simulations there is one atom or several atoms at the center of the cavity. The initial field state is localized on the left and is moving to the right. The atom at the center splits the photon into left and right propagating parts. The spectra of these two parts are measured separately using analyzer atoms and filtered correlation functions. In the first simulation the initial field state is the Gaussian (14) and there is one atom at the center. In the second simulation there are three atoms at the center. In the last simulation there is only one center atom and the field state is a superposition of several random Gaussian states. The units in our simulations are chosen is such a way that $c=\epsilon_0=\mu_0=\hbar=1$.

A. A Gaussian photon and one atom

In the first simulation a Gaussian photon (14) propagates towards the center of the cavity. The length of the cavity is $L=2\pi$. The mode distribution is a Gaussian (14), centered at $k_0=100.0$ with the width $\Gamma_{ph}=4\pi$. The intensity profile is centered at $r_0=2.0$. The initial intensity is shown in Fig. 1.



FIG. 1. The energy density of the photon at $t = 0.0$ and $t = 3.8$. The initial energy density is a Gaussian (14) with parameters $r_0 = 2.0$, $k_0 = 100.0$ and $\Gamma_k = 4\pi$. At $t = 3.8$ there is one peak propagating to the left and two to the right. The initial intensity is split as a results of the interaction with the center atom, marked by a cross. The circles show the positions of the analyzer atoms. The length of the cavity is $L = 2\pi$ and the number of modes is $N_{mode} = 400$.

In the center at $r = \frac{L}{2}$, there is a two level atom which is exactly on resonance with the photon $\omega_0 = 100.0$. The decay constant of the atom is $\Gamma = \pi$, so the linewidth of the atom is narrower than the width of the mode distribution of the field. When the photon reaches the atom, the atom gets a nonzero excitation. Later the atom emits energy back to the field modes via decay to the ground state. The population of the excited state as a function of time is shown in Fig. 2.



FIG. 2. The excitation of the center atom. The photon excites the atom which decays exponentially to the ground state. The decay constant of the atom is $\Gamma = \pi$.

As a result of the interaction, part of the energy is reflected to the left and part is able to pass the atom. The intensity at $t = 3.8$, after the interaction, is shown in Fig. 1. The intensity profile on the right has two peaks. The first peak has propagated to the right without interaction and the second is the result of the atomic decay.

In addition to the center atom there are 600 atoms which detect the spectrum of the radiation which passes them as was explained in section IV. At $r = 1.8$ i.e. left from the center atom there are 200 atoms which measure the spectrum of the reflected radiation. The dipole couplings of the atoms are time dependent in such a way that they are zero when the initial Gaussian photon propagates to the right. At $t = 1.5$ the atomic dipoles get the values determined by (26) and start to detect the spectrum of the radiation. On the right at $r = \frac{L}{2} + 1.0$ there are two groups of analyzer atoms. The first group (200 atoms) measures the spectrum all the time. After the first peak on the right has passed them, its spectrum can be read from the excitations. After the second peak has passed them the atoms give the spectrum of all radiation right from the atom i.e. spectrum of the first and second peaks. Another group at the same position $r = L/2.0 + 1.0$ detects the spectrum of the second peak on the right. The dipole constants of these atoms get nonzero values (26) immediately after the first peak has passed them.

The spectra of the total intensity on the right and left measured using the atoms are shown in Fig. 3.



FIG. 3. The normalized spectra of the radiation on the left and right measured by the analyzer atoms, Fig. 1. The resonant radiation has been reflected to the left and the off-resonant radiation has propagated to the right. The spectrum has been read from the atoms after all radiation has propagated through them. Both spectra have been normalized in such a way that the area under the curves is unity.

On the right the spectrum has a two peak structure. It is interesting that on the right there is no intensity at the resonance frequency of the atom. The center atom has been able to reflect the resonant radiation to the left and only off-resonant radiation passes the atom. The fact that resonant radiation is missing is interesting because the second peak of the intensity profile is the result of the atomic decay.

As was explained earlier, we also measured the spectra of the two peaks on the right separately. The result is shown in Fig. 4.



FIG. 4. The measured spectra of the two peaks on the right (Fig. 1) separately. The spectrum of the first peak is broad because the intensity profile is narrow. The second peak has a typical Lorentzian spectrum of free decay. When the spectrum of both peaks is measured, as in Fig. 3, the second peak creates an stimulated decay type of effect to the detector atoms and the measured spectrum has a dip in the middle. The normalization of the spectra is the same as in Fig. 3

The spectrum of the first peak is quite close to the initial spectrum. The spectrum of the second peak is narrower and is similar to the spectrum of the spontaneous decay. The fact that the spectrum of the second peak is narrower is understandable because the intensity profile is wider and the interaction time with the detector atoms longer. The combined spectrum of the two peaks has a dip in the middle, Fig. 3, whereas the spectra of the peaks measured separately both have a single peak structure. This shows that the spectrum is not additive. The second peak causes a stimulated decay type of effect and the excitations of the detector atoms, which are on resonance with the radiation, get smaller.

Next we calculate the mode spectrum using the filtered correlation functions as was explained in Sec. III. The normally ordered correlation function $\langle : \hat{E}(r_1)\hat{E}(r_2) : \rangle$ at $t = 3.8$ is shown in Fig. 5 (a). The diagonal elements $r_1 = r_2$ give the energy density of the radiation. Off-diagonal elements show the coherence of the radiation. In order to calculate the spectrum of the radiation in the left part of the cavity, a filtered correlation function is used. The part $r_1, r_2 \geq \frac{L}{2}$ is replaced by zero. The filtered correlation function is shown in Fig. 5 (b).



FIG. 5. The correlation function $\langle\Psi| : \hat{E}(r_1)\hat{E}(r_2) : |\Psi\rangle$ of the radiation at $t = 3.8$ (a). The diagonal elements give the energy density of the radiation, Fig. 1. The lower figure (b) shows the filtered correlation function used to determine the mode spectrum on the right.

Similarly in order to get the spectrum of the radiation on the right, the replacement $\langle : \hat{E}(r_1)\hat{E}(r_2) : \rangle = 0$ when $r_1, r_2 \leq \frac{L}{2}$ is used. The mode spectrum is calculated using the formula (20) with a filtered correlation function. The mode spectra obtained are identical to the spectra measured using the analyzer atoms, Fig. 3. We calculated also the mode spectra of the two peaks on the right separately using the appropriate filtered correlation functions. Also these spectra were the same as measured by analyzer atoms, Fig. 4.

B. A Gaussian photon and three atoms

Next we add two more atoms to the center with resonance frequencies and decay constants $\omega_0 = 90.0$, $\Gamma = \pi$ and $\omega_0 = 110.0$, $\Gamma = \pi/4$ respectively. The third atom is the same as earlier, $\omega_0 = 100.0$ and $\Gamma = \pi$. The atom with resonance frequency $\omega_0 = 110.0$ has a smaller decay constant than the other two atoms. The width of the mode spectrum of the Gaussian initial photon is increased to $\Gamma_{ph} = 8\pi$. Because the spectrum is very broad, the energy density profile of the photon is narrow. We have carried out exactly the same spectrum measurement as in the previous simulation for the radiation which passes the atoms and for the reflected radiation. The results are shown in Fig. 6.



FIG. 6. The measured spectra of the radiation on the left and right in the simulation when there are three atoms at the center of the cavity. The radiation on the left has three peaks at resonance frequencies. On the right the broad spectrum has three holes. The width of the peak (or hole) at $\omega_0 = 110.0$ is narrower than the two others. The length of the cavity is $L = 8\pi$ and the number of modes $N_{mode} = 1600$. The normalization of the spectra is the same as in Fig. 3

On the left the spectrum has three peaks at the resonance frequency. On the right the original Gaussian spectrum has three holes at the atomic frequencies. The width of the peak and the hole at $\omega_0 = 110$ is narrower as was expected based on the atomic parameters. The result is qualitatively the same as in the simulation with one atom in the center. The resonant radiation is reflected and the off-resonant radiation is able to pass the atoms. As in the previous simulation we calculated the two spectra using also the filtered correlation functions. The correlation function has a more complicated form than the one in Fig. 5. The result is again identical to the spectra given by the analyzer atoms.

C. A random photon and one atom

In the two previous simulations the initial state of the photon has been of the Gaussian form (14). Next we choose the initial state to be rather exotic. The idea of this simulation is to test the method using a totally different type of field excitation than in the previous simulations. The initial photon is a superposition of ten random Gaussian states. The parameters k_0, σ_k^2 and r_0 in equation (14) are chosen randomly, in such a way that the initial energy density is on the left and propagating to the right. The frequency distribution is centered at $\omega = 100.0$. Again at the center there is one atom which splits the intensity into two parts. The initial intensity profile of the photon is shown in Fig. 7(a).



FIG. 7. The intensity profile (a) and the mode spectrum (b) of the field state which is a sum of ten Gaussian distributions for the photon. The field is propagating to the right. Note that only a part of the cavity is shown. At the center, $r = \frac{L}{2}$, there is one atom with the resonance frequency $\omega_0 = 100.0$. The circles on the left and right are analyzer atom which detect the spectrum. The initial mode spectrum, (b), has a six peak structure as some of the Gaussians are overlapping. The normalization of the spectrum is the same as in Fig. 3

Note that the length of the cavity is now $L = 8\pi$. The field has a four peak structure and it propagates to the right towards the center atom (thick circle). The atom has the resonance frequency $\omega_0 = 100.0$. Circles right and left from the center atom show the positions of the analyzer atoms which detect the spectrum. Figure 7(b) shows the mode spectrum of the initial photon.

As in the earlier simulations the center atom splits the radiation into left and right propagating parts. The spectra of the two parts are shown in Fig. 8.



FIG. 8. The spectra on the left and right measured using analyzer atoms. The initial state of the field is a sum of ten random Gaussian states, Fig. 7. The spectrum has been read from the atoms after all radiation has propagated through them. The normalization of the spectrum is the same as in Fig. 3

Because there is not much intensity which would be at resonance with the center atom, the spectrum on the right is similar to the original mode spectrum. Only a small fraction of the intensity is reflected to the left. Both spectra have several peaks. We calculated the mode spectrum on the left and right separately using filter functions as earlier. Again the mode spectrum gives exactly the same results as detected using analyzer atoms.

VI. CONCLUSION

We have used two level atoms to detect the time dependent and local spectrum of radiation in a one dimensional cavity. The spectrum is read from the excitation probabilities of atoms with very small decay constants. An alternative method to determine the spectrum in this paper has been to calculate the mode coefficients using a filtered second order correlation function. Different filter functions can be used to determine the spectrum at a specific part of the cavity. It might be possible to generalize the approach of filtered correlation functions using wavelet expansions [14–16]. In all cases studied, the spectra determined using these two methods give identical results. In our earlier paper we showed that the analyzer atom spectrum gives the same result for a resonance fluorescence spectrum of a laser driven three level atom as the 'physical spectrum' defined by Eberly and Wódkiewicz. This paper gives further proof that the method really works.

One benefit of the method is that only one time averages of quantum mechanical operators are needed. Thus it is closer to a realistic spectrum measurement than the usual definitions of time dependent spectrum, which typically require two time averages. It can also be used in situations were the usual method to calculate two time averages in the Schrödinger picture, the quantum regression theorem, is known to give incorrect results.

VII. ACKNOWLEDGEMENTS

I want to thank the Academy of Finland for the financial support. Computers of the Center for Scientific Computing (CSC) were used in the simulations. Finally I want to thank V. Bužek, G. Drobný, S. Stenholm and K.-A. Suominen for comments.

---

[1] Page C. H., 1952, *J. Appl. Phys*, **23**, 103.
[2] Lampard D. G., 1954, *J. Appl. Phys*, **25**, 803.
[3] Silverman R. A., 1957, *Proc. I.R.E. (Trans. Inf. Th)*, **3**, 182.
[4] Eberly J. H. and Wódkiewicz K., 1977, *J. Opt. Soc. Am.*, **67**, 1252.
[5] Havukainen M. and Stenholm S., 1998, *J. Mod. Opt.*, **45**(8), 1699.
[6] Lax M., 1963, *Phys. Rev. A*, **129**, 2342.
[7] Ford G. W. and O'Connell R. F., 1996, *Phys. Rev. Lett.*, **77**, 798.
[8] Gardiner C. W., 1993, *Phys. Rev. Lett.*, **70**, 2269.
[9] Carmichael H. J., 1993, *Phys. Rev. Lett.*, **70**, 2273.
[10] Bužek V., 1989, *Czech. J. Phys. B*, **39**, 345.
[11] Bužek V. and Kim M. G., 1997, *J. Korean Phys. Soc.*, **30**, 413.
[12] Bužek V., Drobný G., Kim M. G., Havukainen M. and Knight P. L., 1999, *Phys. Rev. A*, **60**, 582.
[13] Havukainen M., Drobný G., Stenholm S. and Bužek V., 1999, *J. Mod. Opt.*, **46**(9), 1343.
[14] Goswami J. C. and Chan A. K., 1999, *Fundamentals of Wavelets* (Wiley, New York).
[15] Chui C. K., 1992, *An Introduction to Wavelets* (Academic Press, Boston).
[16] Kaiser G., 1994, *A Friendly Guide to Wavelets* (Birkhäuser, Boston).